Author Index

Aburatani, R., see Okamoto, Y 95

Albracht, J. H.
—— and De Wit, M. S.
Analysis of gentamicin in raw material and in pharmaceutical preparations by high-performance liquid chromatography 306

Arentsen-Stasse, A. P., see Van Boekel, M. A. J. S. 267

Armstrong, D. W., see Han, S. M. 256

Asukabe, H., see Oka, H. 417

Bada, J. L., see Gaines, S. M. 219

Barka, G.
—— and Hoffmann, P.
Simultaneous separation of oligomeric and polymeric ethylene glycols (degree of polymerization 1–110) using reversed-phase high-performance liquid chromatography 273

Batley, G. E.
Applications of liquid chromatography with electrochemical detection to the analysis of oil shale process waters 409

Berkowitz, S. A.
—— and Henry, M. P.
Use of high-performance hydrophobic interaction chromatography for the determination of salting-out conditions of proteins 317

Bessalle, R.
—— and Lavie, D.
Semi-quantitative reversed-phase high-performance liquid chromatographic analysis of the ecotypes of *Withania somnifera* chemotype III 195

Bester, A. J., see Victor, T. 339

Bier, M., see Thormann, W. 75

Boekel, M. A. J. S. van, see Van Boekel, M. A. J. S. 267

Bremm, K. D., see Knöller, J. 312

Burcinová, A.
——, Štulík, K. and Pacáková, V.
High-performance liquid chromatography of azobenzene derivatives with spectrophotometric and electrochemical detection 397

Chang, F. C., see Iida, T. 155

Chu, A. H. T.
—— and Langer, S. H.
Determination of the phase ratio by means of reaction kinetics in a liquid chromatographic reactor 11

Churáček, J., see Horna, A. 293

Coenegracht, P. M. J.
——, Van Tuyen, N., Metting, H. J. and Coenegracht-Lamers, P. J. M.
Application of a mixture design technique to the simultaneous optimization of analysis time and resolution in reversed-phase ion-pair liquid chromatography using a minimal resolution plot 351

Coenegracht-Lamers, P. J. M., see Coenegracht, P. M. J. 351

De Wit, M. S., see Albracht, J. H. 306

Dickson, L. C.
—— and Karasek, F. W.
Mechanism of formation of polychlorinated dibenzo-*p*-dioxins produced on municipal incinerator flyash from reactions of chlorinated phenols 127

Dijk, J. A. P. P. van, see Nicolai, T. 286

Dijk, L. van, see Nicolai, T. 286

Djura, P.
—— and Hoskinson, R. M.
^{125}I-Somatostatin analogues: high-performance liquid chromatography profiles and antibody binding properties following three methods of radioiodination 261

Doodokyan, D. L., see Kanavarioti, A. 334

Eckstein, H., see Schott, H. 165

Edberg, S. C., see Torres, A. R. 177

El Tayar, N., see Tsantili-Kakoulidou, A. 33

Erdelmeier, C. A. J.
——, Kinghorn, A. D. and Farnsworth, N. R.
On-plate injection in the preparative separation of alkaloids and glycosides using overpressured layer chromatography 345

Evans, M. B.
—— and Haken, J. K.
Dispersion and selectivity indices of the halogenated derivatives of cyclohexane, benzene and anisole 240

Farnsworth, N. R., see Erdelmeier, C. A. J. 345

Fonck, C., see Hughes, G. J. 327

Frutiger, S., see Hughes, G. J. 327

Gaines, S. M.
—— and Bada, J. L.
Reversed-phase high-performance liquid chromatographic separation of aspartame diastereomeric decomposition products 219

Gale, R. W., see Yonker, C. R. 433

Gether, U., see O'Hare, M. M. T. 379

Gharfeh, S. G.
Determination of the hindered amine additive Chimassorb 944 in polyethylene by high-performance liquid chromatography 211
Goto, J., see Iida, T. 155
Haddad, P. R., see Jackson, P. E. 65
Hagenaars, A., see Wong, D. 279
Haikala, V.
Rapid high-performance liquid chromatography determination of noscapine hydrogen embonate 299
Haken, J. K., see Evans, M. B. 240
Han, S. M.
—— and Armstrong, D. W.
Use of microcolumn liquid chromatography with a chiral stationary phase for the separation of low-resolution enantiomers 256
Harada, K.-I., see Oka, H. 417
Hatada, K., see Okamoto, Y. 95
Henry, M. P., see Berkowitz, S. A. 317
Hicks, K. B.
—— and Sondey, S. M.
Preparative high-performance liquid chromatography of malto-oligosaccharides 183
Hoffmann, P., see Barka, G. 273
Horna, A.
—— and Churáček, J.
Chromatography of monomers. VII. Determination of residual ethyl acrylate and methyl methacrylate in polymer emulsion: effect of precipitated copolymer 293
Hoskinson, R. M., see Djura, P. 261
Hostettmann, K., see Schaufelberger, D. 450
Hughes, G. J.
——, Frutiger, S. and Fonck, C.
Quantitative high-performance liquid chromatographic analysis of Dabsyl-amino acids within 14 min 327
Iida, T.
——, Momose, T., Tamura, T., Matsumoto, T., Goto, J., Nambara, T. and Chang, F.C.
Capillary gas chromatography of hydroxylated bile acid stereoisomers of the *allo* and normal series 155
Ikai, Y., see Oka, H. 417
Jackson, P. E.
—— and Haddad, P. R.
Studies on sample preconcentration in ion chromatography. V. Effect of the ion-exchange capacity of the concentrator column 65
Jalsovszky, I.
——, Szókán, G., Ruff, F, and Kucsman, Á.
High-performance liquid chromatography of cyclic sulphilimine and sulphoxide diastereoisomers 439
Jandik, P., see Wong, D. 279
Jefferies, T. M., see Patel, H. B. 21
Johansen, M., see Larsen, C. 227
Jones, W. R., see Wong, D. 279
Jordaan, A. M., see Victor, T. 339
Kanavarioti, A.
—— and Doodokyan, D. L.
High-performance liquid chromatographic method using a C_{18} column for the simultaneous separation of the products of decomposition and oligomerization of guanosine 5'-phospho-2-methylimidazolide 334
Kaneshima, H., see Kanetoshi, A. 139
Kanetoshi, A.
——, Ogawa, H., Katsura, E. and Kaneshima, H.
Chlorination of Irgasan DP300 and formation of dioxins from its chlorinated derivatives 139
Karasek, F. W., see Dickson, L. C. 127
Katsura, E., see Kanetoshi, A. 139
Kawamura, N., see Oka, H. 417
Kennedy, J. H.
—— and Olsen, B. A.
Investigation of perchlorate, phosphate and ion-pairing eluent modifiers for the separation of cephalosporin epimers 369
Kinghorn, A. D., see Erdelmeier, C. A. J. 345
Kitaura, F., see Miyake, K. 47
Knöller, J.
——, Schönfeld, W., Bremm, K. D. and König, W.
In vitro stability of cefixime (FK-027) in serum, urine and buffer 312
König, W., see Knöller, J. 312
Kovář, J.
——, Pavlicová, Z., Zemek, J. and Kuniak, L.
Chromatofocusing of cellulolytic enzymes produced by *Trichoderma viride, Trichoderma reesei, Aspergillus niger* and *Pleurotus ostreatus* 322
Kucsman, Á., see Jalsovszky, I. 439
Kuniak, L., see Kovář, J. 322
Langer, S. H., see Chu, A. H. T. 11
Larsen, C.
—— and Johansen, M.
Macromolecular prodrugs. V. Simultaneous determination of high-molecular-weight dextran metronidazole monosuccinate ester prodrugs and the hydrolysis products metronidazole and the corresponding monosuccinate ester on nucleosil diol 227
Lavie, D., see Bessalle, R. 195
Lentzen, H.
—— and Simon, R.
Reversed-phase high-performance liquid chromatography of tyrosine-containing metabolites of enkephalins using octanesulphonic acid as ion-pairing agent and radioactivity detection 444

Leyendecker, D., see Schmitz, F. P. 245
Lochner, A., see Victor, T. 339
Matsumoto, T., see Iida, T. 155
Metting, H. J., see Coenegracht, P. M. J. 351
Michaud, J.-P., see Thormann, W. 75
Miyake, K.
——, Kitaura, F., Mizuno, N. and Terada, H.
Phosphatidylcholine-coated silica as a useful stationary phase for high-performance liquid chromatography determination of partition coefficients between octanol and water 47
Mizuno, N., see Miyake, K. 47
Momose, T., see Iida, T. 155
Morecombe, D. J.
High-efficiency preparative-scale reversed-phase high-performance liquid chromatographic purification of ^{14}C-labelled antibiotics 389
Mori, Y., see Oka, H. 417
Mosher, R. A., see Thormann, W. 75
Nambara, T., see Iida, T. 155
Nicolai, T.
——, Van Dijk, L., Van Dijk, J. A. P. P. and Smit, J. A. M.
Molar mass characterization of DNA fragments by gel permeation chromatography using a low-angle laser light-scattering detector 286
Nielsen, H. V., see O'Hare, M. M. T. 379
Ogawa, H., see Kanetoshi, A. 139
O'Hare, M. M. T.
——, Tortora, O., Gether, U., Nielsen, H. V. and Schwartz, T. W.
High-performance liquid chromatography of phenylthiocarbamyl derivatives of amino acids and side-chain derivatized amino acids 379
Ohya, K., see Takasu, A. 251
Oka, H.
——, Ikai, Y., Kawamura, N., Uno, K., Yamada, M., Harada, K.-I., Uchiyama, M., Asukabe, H., Mori, Y. and Suzuki, M.
Improvement of chemical analysis of antibiotics. IX. A simple method for residual tetracyclines analysis in honey using a tandem cartridge clean-up system 417
Okamoto, Y.
——, Aburatani, R. and Hatada, K.
Chromatographic chiral resolution. XIV. Cellulose tribenzoate derivatives as chiral stationary phases for high-performance liquid chromatography 95
Olsen, B. A., see Kennedy, J. H. 369
Pacáková, V., see Burcinová, A. 397
Patel, H. B.
—— and Jefferies, T. M.
Eluotropic strength of solvents. Prediction and use in reversed-phase high-performance liquid chromatography 21
Patton, E. V., see Wonnacott, D. M. 103
—— and Wonnacott, D. M.
Characterization of polymeric aminopropyl derivatized silica support for the size-exclusion chromatography of cationic polymers 115
Pavlicová, Z., see Kovář, J. 322
Perry, J. A.
——, Rateike, J. D. and Szczerba T. J.
Eluting trace components before major constituents. I. Sensitivity enhancement in analytical determinations of optical purity 57
Peterson, E. A., see Torres, A. R. 177
Prud'homme, R. K., see Sugarman, J. H. 236
Rateike, J. D., see Perry, J. A. 57
Ruff, F., see Jalsovszky, I. 439
Schaufelberger, D.
—— and Hostettmann, K.
High-performance liquid chromatographic analysis of secoiridoid and flavone glycosides in closely related *Gentiana* species 450
Schmitz, F. P.
——, Leyendecker, D. and Leyendecker, D.
Permanent "on column" modification of a silica stationary phase with 1,4-dioxane 245
Schönfeld, W., see Knöller, J. 312
Schott, H.
——, Semmler, R. and Eckstein, H.
Column chromatographic purification of guanylate-rich synthetic oligodeoxyribonucleotides 165
Schwartz, T. W., see O'Hare, M. M. T. 379
Semmler, R., see Schott, H. 165
Simon, R., see Lentzen, H. 444
Sipponen, K. B.
Detector for organophosphorus compounds in liquid chromatography based on the cholinesterase inhibition reaction 87
Skorkowski, E. F.
—— and Storey, K. B.
Affinity chromatography on 2',5'-ADP-Sepharose 4B for purification of malic enzyme from crustacean muscle 427
Smit, J. A. M., see Nicolai, T. 286
Smith, R. D., see Yonker, C. R. 433
Sondey, S. M., see Hicks, K. B. 183
Storey, K. B., see Skorkowski, E. F. 427
Štulík, K., see Burcinová, A. 397

Sugarman, J. H.
—— and Prud'homme R. K.
Analysis of radius variations in microcapillaries using the velocity of a viscous polymer plug and fluorescence photobleaching velocity detection 236
Suzuki, M., see Oka, H. 417
Szczerba, T. J., see Perry, J. A. 57
Szókán, G., see Jalsovszky, I. 439
Takasu, A.
—— and Ohya, K.
Separation and determination of the enantiomers of pantolactone by gas–liquid chromatography 251
Takeuchi, T.
—— and Yeung, E. S.
Signal enhancement in on-column fluorometric detection in open-tubular capillary liquid chromatography 3
Tamura, T., see Iida, T. 155
Tayar, N. El, see Tsantili-Kakoulidou, A. 33
Terada, H., see Miyake, K. 47
Testa, B., see Tsantili-Kakoulidou, A. 33
Thormann, W.
——, Tsai, A., Michaud, J.-P., Mosher, R. A. and Bier, M.
Capillary isoelectric focusing. Effects of capillary geometry, voltage gradient and addition of linear polymer 75
Torres, A. R.
——, Edberg, S. C. and Peterson, E. A.
Preparative high-performance liquid chromatography of proteins on an anion exchanger using unfractionated carboxymethyldextran displacers 177
Tortora, O., see O'Hare, M. M. T. 379
Tsai, A., see Thormann, W. 75
Tsantili-Kakoulidou, A.
——, El Tayar, N., Van de Waterbeemd, H. and Testa, B.
Structural effects in the lipophilicity of di- and polysubstituded benzenes as measured by reversed-phase high-performance liquid chromatography 33
Tuyen, N. van, see Coenegracht, P. M. J. 351
Uchiyama, M., see Oka, H. 417
Uno, K., see Oka, H. 417
Van Boekel, M. A. J. S.
—— and Arentsen-Stasse, A. P.
Determination of aromatic biogenic amines and their precursors in cheese by high-performance liquid chromatography 267
Van de Waterbeemd, H., see Tsantili-Kakoulidou, A. 33
Van Dijk, J. A. P. P., see Nicolai, T. 286
Van Dijk, L., see Nicolai, T. 286
Van Tuyen, N., see Coenegracht, P. M. J. 351
Victor, T.
——, Jordaan, A. M., Bester, A. J. and Lochner, A.
A sensitive and rapid method for separating adenine nucleotides and creatine phosphate by ion-pair reversed-phase high-performance liquid chromatography 339
Waterbeemd, H. Van de, see Tsantilli-Kakoulidou, A. 33
Wit, M. S. de, see Albracht, J. H. 306
Wong, D.
——, Jandik, P., Jones, W. R. and Hagenaars, A.
Ion chromatography of polyphosphonates with direct refractive index detection 279
Wonnacott, D. M., see Patton, E. V. 115
—— and Patton, E. V.
Hydrolytic stability of aminopropyl stationary phases used in the size-exclusion chromatography of cationic polymers 103
Yamada, M., see Oka, H. 417
Yeung, E. S., see Takeuchi, T. 3
Yonker, C. R.
——, Gale, R. W. and Smith, R. D.
Effect of solute concentration on retention in supercritical fluid chromatography 433
Zemek, J., see Kovář, J. 322